

# ARE FEEDING TRAITS AND HABITAT RESPONSIBLE OF MICROPLASTICS INGESTION IN FISH, CRUSTACEANS AND ELASMOBRANCHS AT THE WESTERN MEDITERRANEAN?

Salud Deudero <sup>1\*</sup> and Carme Alomar Mascaró <sup>1</sup>

<sup>1</sup> Instituto Español de Oceanografía (IEO) Centro Oceanográfico de Baleares - salud.deudero@ba.ieo.es

## Abstract

Marine litter loads are increasing worldwide and impacts and effects on marine ecosystems and their inhabitants are still unknown [1,2]. Whereas interaction effects of macrolitter, especially on species as sea turtles and marine mammals has been more investigated, the microscopic fraction has been less addressed. Therefore, several key species of fish, crustaceans and elasmobranchs have been studied to assess microplastics ingestion in the Western Mediterranean. Mean ingested microplastics (MPs) ranged up to 2.3 MPs/ind indicating a threat of this man made contaminant on species which are commercialised.

**Keywords** *South-Western Mediterranean, Plastics, Pollution*

## Introduction

The small size of MPs facilitates organisms' intake compared to macroplastics and widespread ecological impacts are expected from MPs. Quantities of MPs have already been reported in shallow marine coastal areas of the Balearic Islands [3]. Therefore, MPs loads in the marine environment are available for organisms with different feeding strategies which can ingest them randomly or selectively having implications along the food web.

## Material and Methods

Fishes, crustaceans and elasmobranchs exhibiting different trophic strategies and trophic levels have been taken as case study species to evaluate microplastics ingestion (<5mm plastic fragments). Stomach contents analyses of the selected species have been conducted to quantify mean number of MPs/ind [4,5]. Species have been classified into several feeding habits: browsing on substrate, hunting for macrofauna (predators), selective plankton feeding and variable, and analysed according to their environment: bathydemersal, bathypelagic, and demersal and pelagic-neritic.

## Results and Discussion

Different results reveal MPs ingestion of commercial fish and crustacean species *Boops boops*, *Mullus surmuletus*, *Aristeus antennatus*, among others (Fig 1). Most species ingested less than 0.5 MPs/ind. However, *B. boops* is the most affected species exhibiting ingestion rates up to 2.3 MPs/ind being mainly filament type MPs. Results indicate that combined functional traits: biology, autoecology, ethology could be responsible of microplastics ingestion at the studied species. Microplastics are menacing functional diversity through differential ingestion linked to organisms' trophic strategies.

## Acknowledgements

We appreciate T. Box (Consell d'Eivissa i Formentera), A. de Mesa, MEDITS and COB for providing species samples.

## References

- 1 - Deudero, S and Alomar, C 2015 Mediterranean marine biodiversity under threat: reviewing influence of marine litter on species. *Marine Pollution Bulletin*, 98: 58-68
- 2 - Deudero, S and Alomar, C 2014 Revising interactions of plastics with marine biota: evidence from the Mediterranean in CIESM 2014, *Marine litter in the Mediterranean and Black Seas*. CIESM Workshop Monograph n° 46 [F. Briand, ed.], 180 p., CIESM Publisher, Monaco
- 3 - Alomar, C, Estarellas, F., Deudero, S 2016 Microplastics in the Mediterranean sea: Deposition in coastal shallow sediments, spatial variation and preferential grain size. *Marine Environmental Research*, 115: 1-10. doi:10.1016/j.marenvres.2016.01.005
- 4 - Deudero, S., Nadal, M.A., Estarellas, F., Alomar, C 2014 Microplastic exposure in pelagic fishes and holothurians: a Mediterranean case study. 2nd International Ocean Research Conference Barcelona 17-21 November 2014
- 5 - Frank, Aida 2015 Microplastic ingestion in elasmobranchs from the Western Mediterranean. University of the Balearic Islands, Final Degree Thesis in Biology. September 2015




Fig 1 Mean microplastics (MPs) per individual according to inhabiting environment