



Gene expression of *Solea senegalensis* Kaup, 1858 fed with the synbiotic composed by sodic alginate and the probiotic *Shewanella putrefaciens* Pdp11


Danielle De Carla Dias ^{1,2}, Sara Vidal ², Silvana Tapia-Paniagua ², Carmen Lobo ³, Inés García de la Banda ³, María Carmen Balebona ², Miguel Ángel Moriñigo ².
 1. Research Center in Aquaculture, Fisheries Institute, São Paulo, Brazil. 2. Microbiology, University of Málaga, Málaga. 3. Spanish Institute of Oceanography, Santander, Spain

Introduction

The culture of the Senegalese sole (*Solea senegalensis*, Kaup 1858) is a promising industry in the Mediterranean countries. This species is very attractive for aquaculture, but the problems for controlling infectious diseases can seriously constrain its production. The use of antibiotics can control most diseases, but they also have adverse effects such as accumulation in the tissue, immunosuppression, and development of antibiotic resistant bacteria. The use of probiotics and prebiotics seems to be a favorable alternative to antibiotics.

Previous studies showed that probiotic *Shewanella putrefaciens* Pdp11 has the ability to control the infection by several pathogens in *S. senegalensis*. The administration of the probiotic increased fish growth, enhanced resistance to diseases and induced changes in the intestinal microbiota. On the other hand, prebiotics are non-digestible food ingredients that stimulate the growth and/or activity of bacteria in the digestive system leading to benefits for the host. Nowadays, some studies are focused in the co-administration effects of probiotic and prebiotic as a synbiotic. The present work investigates the effect of *S. putrefaciens* Pdp11 (as probiotic) administered together with alginate (as prebiotic), on the immune system of cultured Senegalese sole.

Material and methods




Target genes				
Protein	Gene	Primers (5'-3')	Amplicon (pb)	References
C3 component	C3	F: ACCTTAGACTGCCCTACTCTGCTGTCCGTG R: GCACTGCACACATCATCCGTCTCAGAC	127	Prieto-Álamo <i>et al.</i> , 2009
C7 component	C7	F: GGCACACACTATCTGTCGCAGGGCTC R: GGCGAACGCCTGATGGTTAACTCCAG	78	Prieto-Álamo <i>et al.</i> , 2009
Lysozyme G	LYSG	F: ACTGCTCGCGGTGAATGGGACA R: CCTGAAAATTTATTACGGATTCCGCCAATG	95	Salas-Leiton <i>et al.</i> , 2010
Glutathion peroxidase	GPXI	F: GATTCGTCCAAACTTCCTGCTA R: GCTCCCAGAACGCCTGTTG	212	Teles <i>et al.</i> , 2011
NADPH oxidase	CYBB	F: CATCGCCACCTGTTAACT R: GTATGACCTGCGGATGACCT	250	Teles <i>et al.</i> , 2011

Housekeeping genes			
Gene	Primers (5'-3')	Amplicon (pb)	References
ACTB2	F: AATCGTGACCTCTGCTTCCCCTGT R: TCTGGCACCCCATGTTACCCCATC	113	Infante <i>et al.</i> , 2008
RPS4	F: GTGAAGAAGCTCCTTGTCGGCACCA R: AGGGGGTCGGGTAGCGGATG	83	Infante <i>et al.</i> , 2008

Discussion and conclusions

After 10 days feeding with the synbiotic, gene expression showed a significant upregulation in comparison to control fish in all studied genes. These results can demonstrate that the synbiotic can alter the expression of genes related with immune system functions. The immuno-stimulation produced by the synbiotic may provide an additional overprotection against pathogens or stressors processes. However, the permanent application of the synbiotic and the continuous immuno-stimulation could be harmful to the fish. Therefore, more studies are necessary to see how would affect to fish if we still adding the synbiotic and if the synbiotic could give protection against experimental infections.

Results



References

- Infante C, Matsuoka P, Asensio, Cañavate JP, Reith, Manchado M. (2008). Selection of housekeeping genes for gene expression studies in larvae from flatfish using real-time PCR. *BMC Molecular Biology* 9:28, *Fish & Shellfish Immunology* 28, 296–302.
- Prieto-Álamo MJ, Abril N, Osuna-Jiménez I, Pueyo C. (2009). *Solea senegalensis* genes responding to lipopolysaccharide and copper sulphate challenges: Large-scale identification by suppression subtractive hybridization and absolute quantification of transcriptional profiles by real-time RT-PCR. *Aquatic Toxicology* 91, 312-319.
- Salas-Leiton E, Anguís V, Rodríguez-Rúa A, Cañavate JP. (2010). Stocking homogeneous size groups does not improve growth performance of Senegalese sole (*Solea senegalensis*, Kaup 1858) juveniles: Individual growth related to fish size. *Aquacultural Engineering* 43, 108-113.
- Teles M, Mackenzie S, Boltaña S, Callol A. (2011). Gene expression and TNF-alpha secretion profile in rainbow trout macrophages following exposures to copper and bacterial lipopolysaccharide. *Fish & Shellfish Immunology* 30 (1), 340–346.